arXiv:1211.2909v1 [nlin.SI] 13 Nov 2012

BOSONIZATION AND VERTEX OPERATOR OF SUPERSYMMETRY $U_q(\widehat{sl}(N|1))$ FOR LEVEL k

November 14, 2012

TAKEO KOJIMA

Faculty of Engineering, Yamagata University, Jonan 4-3-16, Yonezawa 992-8510, Japan
kojima@yz.yamagata-u.ac.jp

Abstract

We construct a bosonization of the quantum superalgebra $U_q(\hat{sl}(N|1))$ for an arbitrary level k. We construct the screening that commutes with the quantum superalgebra for an arbitrary level $k \neq -N+1$. We propose a bosonization of the vertex operator that gives the intertwiner among the Wakimoto realization and the typical representation.

1 Introduction

There have been many developments in exactly solvable models. Various methods were invented to solve models. The bosonization provides a powerful method to study exactly solvable models. We review recent developments in bosonizations of the $U_q(\widehat{sl}(N|1))$ [7, 8, 10]. The trace of our bosonizations of the vertex operators gives the correlation function of the higher level(spin) k and rank N generalization of the supersymmetric t-J model ($U_q(\widehat{sl}(2|1))$ for level $k=1$) [1].

2 Bosonization of $U_q(\widehat{sl}(N|1))$

We recall Drinfeld generators of the superalgebra $U_q(\widehat{sl}(N|1))$ [3]. We fix a complex number $q \neq 0, |q| < 1$. We use the notations $[x,y] = xy - yx$, $\{x,y\} = xy + yx$, $[a]_q = \frac{q^a - q^{-a}}{q - q^{-1}}$. The Cartan matrix $(A_{i,j})_{0 \leq i,j \leq N}$

of the affine Lie algebra $\widehat{sl}(N|1)$ is given by

$$A_{i,j}=(\nu_i+\nu_{i+1})\delta_{i,j}-\nu_i\delta_{i,j+1}-\nu_{i+1}\delta_{i+1,j}.$$

Here we set $\nu_1=\cdots=\nu_N=+,\nu_{N+1}=\nu_0=-$. The Drinfeld generators of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ are $x_{i,m}^{\pm},\ h_{i,m},\ c,\ (1\leq i\leq N, m\in\mathbf{Z})$. Defining relations are

$$\begin{aligned}
&c:\text{central},\ [h_i,h_{j,m}]=0,\\
&[a_{i,m},h_{j,n}]=\frac{[A_{i,j}m]_q[cm]_q}{m}q^{-c|m|}\delta_{m+n,0}\ \ (m,n\neq 0),\ [h_i,x_j^{\pm}(z)]=\pm A_{i,j}x_j^{\pm}(z),\\
&[h_{i,m},x_j^{+}(z)]=\frac{[A_{i,j}m]_q}{m}q^{-c|m|}z^m x_j^{+}(z),\ \ [h_{i,m},x_j^{-}(z)]=-\frac{[A_{i,j}m]_q}{m}z^m x_j^{-}(z)\ \ (m\neq 0),\\
&(z_1-q^{\pm A_{i,j}}z_2)x_i^{\pm}(z_1)x_j^{\pm}(z_2)=(q^{\pm A_{j,i}}z_1-z_2)x_j^{\pm}(z_2)x_i^{\pm}(z_1)\quad \text{for } |A_{i,j}|\neq 0,\\
&x_i^{\pm}(z_1)x_j^{\pm}(z_2)=x_j^{\pm}(z_2)x_i^{\pm}(z_1)\quad \text{for } |A_{i,j}|=0,(i,j)\neq(N,N),\quad \{x_N^{\pm}(z_1),x_N^{\pm}(z_2)\}=0,\\
&[x_i^{+}(z_1),x_j^{-}(z_2)]=\frac{\delta_{i,j}}{(q-q^{-1})z_1z_2}\left(\delta(q^{-c}z_1/z_2)\psi_i^{+}(q^{\frac{c}{2}}z_2)-\delta(q^{c}z_1/z_2)\psi_i^{-}(q^{-\frac{c}{2}}z_2)\right),\\
&\quad \text{for }\ (i,j)\neq(N,N),\\
&\{x_N^{+}(z_1),x_N^{-}(z_2)\}=\frac{1}{(q-q^{-1})z_1z_2}\left(\delta(q^{-c}z_1/z_2)\psi_N^{+}(q^{\frac{c}{2}}z_2)-\delta(q^{c}z_1/z_2)\psi_N^{-}(q^{-\frac{c}{2}}z_2)\right),\\
&\left(x_i^{\pm}(z_1)x_i^{\pm}(z_2)x_j^{\pm}(z)-(q+q^{-1})x_i^{\pm}(z_1)x_j^{\pm}(z)x_i^{\pm}(z_2)+x_j^{\pm}(z)x_i^{\pm}(z_1)x_i^{\pm}(z_2)\right)\\
&+(z_1\leftrightarrow z_2)=0\quad \text{for } |A_{i,j}|=1,\ i\neq N,
\end{aligned}$$

where we have used $\delta(z)=\sum_{m\in\mathbf{Z}}z^m$. Here we have used the generating function $x_j^{\pm}(z)=\sum_{m\in\mathbf{Z}}x_{j,m}^{\pm}z^{-m-1}$, $\psi_i^{\pm}(q^{\pm\frac{c}{2}}z)=q^{\pm h_i}e^{\pm(q-q^{-1})\sum_{m>0}h_{i,\pm m}z^{\mp m}}$ and the abbreviation $h_i=h_{i,0}$.

We construct bosonizations of superalgebra $U_q(\hat{sl}(N|1))$ for an arbitrary level [7]. We fix the level $c=k\in\mathbf{C}$. We introduce the bosons and the zero-mode operators a_m^j,Q_a^j $(m\in\mathbf{Z},1\leq j\leq N)$, $b_m^{i,j},Q_b^{i,j}$ $(m\in\mathbf{Z},1\leq i<j\leq N+1)$, $c_m^{i,j},Q_c^{i,j}$ $(m\in\mathbf{Z},1\leq i<j\leq N)$ which satisfy

$$\begin{aligned}
&[a_m^i,a_n^j]=\frac{[(k+N-1)m]_q[A_{i,j}m]_q}{m}\delta_{m+n,0}\quad (m,n\neq 0),\quad [a_0^i,Q_a^j]=(k+N-1)A_{i,j},\\
&[b_m^{i,j},b_n^{i',j'}]=-\nu_i\nu_j\frac{[m]_q^2}{m}\delta_{i,i'}\delta_{j,j'}\delta_{m+n,0}\quad (m,n\neq 0),\quad [b_0^{i,j},Q_b^{i',j'}]=-\nu_i\nu_j\delta_{i,i'}\delta_{j,j'},\\
&[c_m^{i,j},c_n^{i',j'}]=\frac{[m]_q^2}{m}\delta_{i,i'}\delta_{j,j'}\delta_{m+n,0}\quad (m,n\neq 0),\quad [c_0^{i,j},Q_c^{i',j'}]=\delta_{i,i'}\delta_{j,j'},\\
&[Q_b^{i,j},Q_b^{i',j'}]=\delta_{j,N+1}\delta_{j',N+1}\pi\sqrt{-1}\qquad ((i,j)\neq(i',j')).
\end{aligned}$$

Other commutation relations are zero. In what follows we use the standard symbol of the normal orderings :: and use the following abbreviations $b^{i,j}(z),c^{i,j}(z),b_{\pm}^{i,j}(z),a_{\pm}^j(z)$ and $\left(\frac{\gamma_1}{\beta_1}\frac{\gamma_2}{\beta_2}\cdots\frac{\gamma_r}{\beta_r}\ a^i\right)(z|\alpha)$ given by

$$\begin{aligned}
b^{i,j}(z)&=-\sum_{m\neq 0}\frac{b_m^{i,j}}{[m]_q}z^{-m}+Q_b^{i,j}+b_0^{i,j}\log z,\ c^{i,j}(z)=-\sum_{m\neq 0}\frac{c_m^{i,j}}{[m]_q}z^{-m}+Q_c^{i,j}+c_0^{i,j}\log z,\\
b_{\pm}^{i,j}(z)&=\pm(q-q^{-1})\sum_{\pm m>0}b_m^{i,j}z^{-m}\pm b_0^{i,j}\log q,\ a_{\pm}^j(z)=\pm(q-q^{-1})\sum_{\pm m>0}a_m^j z^{-m}\pm a_0^j\log q,\\
\left(\frac{\gamma_1}{\beta_1}\frac{\gamma_2}{\beta_2}\cdots\frac{\gamma_r}{\beta_r}\ a^i\right)(z|\alpha)&=-\sum_{m\neq 0}\frac{[\gamma_1 m]_q\cdots[\gamma_r m]_q}{[\beta_1 m]_q\cdots[\beta_r m]_q}\frac{a_m^i}{[m]_q}q^{-\alpha|m|}z^{-m}+\frac{\gamma_1\cdots\gamma_r}{\beta_1\cdots\beta_r}(Q_a^i+a_0^i\log z).
\end{aligned}$$

The generating functions $x_i^\pm(z)$, $\psi_i^\pm(z)$, $(1 \leq i \leq N)$ of $U_q(\widehat{sl}(N|1))$ for an arbitrary level k are realized by the bosonic operators as follows. This is main result of [7].

$$
\begin{aligned}
x_i^+(z) &= \frac{1}{(q-q^{-1})z} : \sum_{j=1}^{i} e^{(b+c)^{j,i}(q^{j-1}z)+\sum_{l=1}^{j-1}(b_+^{l,i+1}(q^{l-1}z)-b_+^{l,i}(q^l z))} \times \\
&\times \left\{ e^{b_+^{j,i+1}(q^{j-1}z)-(b+c)^{j,i+1}(q^j z)} - e^{b_-^{j,i+1}(q^{j-1}z)-(b+c)^{j,i+1}(q^{j-2}z)} \right\} :, \\
x_N^+(z) &= : \sum_{j=1}^{N} e^{(b+c)^{j,N}(q^{j-1}z)+b^{j,N+1}(q^{j-1}z)-\sum_{l=1}^{j-1}(b_+^{l,N+1}(q^l z)+b_+^{l,N}(q^l z))} :, \\
x_i^-(z) &= q^{k+N-1} : e^{a_+^i(q^{\frac{k+N-1}{2}}z)-b^{i,N+1}(q^{k+N-1}z)-b_+^{i+1,N+1}(q^{k+N-1}z)+b^{i+1,N+1}(q^{k+N}z)} : \\
&+ \frac{1}{(q-q^{-1})z} : \left\{ \sum_{j=1}^{i-1} e^{a_-^i(q^{-\frac{k+N-1}{2}}z)+(b+c)^{j,i+1}(q^{-k-j}z)+b_-^{i,n+1}(q^{-k-n}z)-b_-^{i+1,n+1}(q^{-k-n+1}z)} \right. \\
&\times e^{\sum_{l=j+1}^{i}(b_-^{l,i+1}(q^{-k-l+1}z)-b_-^{l,i}(q^{-k-l}z))+\sum_{l=i+1}^{N}(b_-^{i,l}(q^{-k-l}z)-b_-^{i+1,l}(q^{-k-l+1}z))} \\
&\times \left(e^{-b_-^{j,i}(q^{-k-j}z)-(b+c)^{j,i}(q^{-k-j+1}z)} - e^{-b_+^{j,i}(q^{-k-j}z)-(b+c)^{j,i}(q^{-k-j-1}z)} \right) \\
&+ e^{a_-^i(q^{-\frac{k+N-1}{2}}z)+(b+c)^{i,i+1}(q^{-k-i}z)+\sum_{l=i+1}^{N}(b_-^{i,l}(q^{-k-l}z)-b_-^{i+1,l}(q^{-k-l+1}z))+b_-^{i,N+1}(q^{-k-N}z)-b_-^{i+1,N+1}(q^{-k-N+1}z)} \\
&- e^{a_+^i(q^{\frac{k+N-1}{2}}z)+(b+c)^{i,i+1}(q^{k+i}z)+\sum_{l=i+1}^{N}(b_+^{i,l}(q^{k+l}z)-b_+^{i+1,l}(q^{k+l-1}z))+b_+^{i,N+1}(q^{k+N}z)-b_+^{i+1,N+1}(q^{k+N-1}z)} \\
&- \sum_{j=i+1}^{N-1} e^{a_+^i(q^{\frac{k+N-1}{2}}z)+(b+c)^{i,j+1}(q^{k+j}z)+b_+^{i,N+1}(q^{k+N}z)-b_+^{i+1,N+1}(q^{k+N-1}z)+\sum_{l=j+1}^{N}(b_+^{i,l}(q^{k+l}z)-b_+^{i+1,l}(q^{k+l-1}z))} \\
&\times \left. \left(e^{b_+^{i+1,j+1}(q^{k+j}z)-(b+c)^{i+1,j+1}(q^{k+j+1}z)} - e^{b_-^{i+1,j+1}(q^{k+j}z)-(b+c)^{i+1,j+1}(q^{k+j-1}z)} \right) \right\} : . \\
x_N^-(z) &= \frac{1}{(q-q^{-1})z} : \left\{ \sum_{j=1}^{N-1} e^{a_-^N(q^{-\frac{k+N-1}{2}}z)-b_+^{j,N+1}(q^{-k-j}z)-b^{j,N+1}(q^{-k-j-1}z)-\sum_{l=j+1}^{N-1}(b_-^{l,N}(q^{-k-l}z)+b_-^{l,N+1}(q^{-k-l}z))} \right. \\
&\times \left. q^{j-1} \left(e^{-b_+^{j,N}(q^{-k-j}z)-(b+c)^{j,N}(q^{-k-j-1}z)} - e^{-b_-^{j,N}(q^{-k-j}z)-(b+c)^{j,N}(q^{-k-j+1}z)} \right) \right\} : \\
&+ q^{N-1} : \left(e^{a_+^N(q^{\frac{k+N-1}{2}}z)-b^{N,N+1}(q^{k+N-1}z)} - e^{a_-^N(q^{-\frac{k+N-1}{2}}z)-b^{N,N+1}(q^{-k-N+1}z)} \right) : .
\end{aligned}
$$

$$
\begin{aligned}
\psi_i^\pm(q^{\pm\frac{k}{2}}z) &= e^{a_\pm^i(q^{\pm\frac{k+N-1}{2}}z)+\sum_{l=1}^{i}(b_\pm^{l,i+1}(q^{\pm(l+k-1)}z)-b_\pm^{l,i}(q^{\pm(l+k)}z)} \\
&\quad \times e^{\sum_{l=i+1}^{N}(b_\pm^{i,l}(q^{\pm(k+l)}z)-b_\pm^{i-1,l}(q^{\pm(k+l-1)}z)+b_\pm^{i,N+1}(q^{\pm(k+N)}z)-b_\pm^{i+1,N+1}(q^{\pm(k+N-1)}z)}, \\
\psi_N^\pm(q^{\pm\frac{k}{2}}z) &= e^{a_\pm^N(q^{\pm\frac{k+N-1}{2}}z)-\sum_{l=1}^{N-1}(b_\pm^{l,N}(q^{\pm(k+l)}z)+b_\pm^{l,N+1}(q^{\pm(k+l)}z))}.
\end{aligned}
$$

3 Vertex Operator

We construct the screening $\mathcal{Q}_i$ that commutes with the quantum superalgebra $U_q(\widehat{sl}(N|1))$ for an arbitrary level $k \neq -N+1$ [10]. The Jackson integral with parameter $p \in \mathbf{C}$ $(|p| < 1)$ and $s \in \mathbf{C}^*$ is defined by $\int_0^{s\infty} f(z) d_p z = s(1-p) \sum_{m \in \mathbf{Z}} f(sp^m) p^m$. We introduce the screening operators $\mathcal{Q}_i$ $(1 \leq i \leq N)$ by using the Jackson integral. This is one of main result of [10].

$$
\mathcal{Q}_i = \int_0^{s\infty} : e^{-\left(\frac{1}{k+N-1}a^i\right)\left(z\middle|\frac{k+N-1}{2}\right)} \widetilde{S}_i(z) : d_p z, \quad (p = q^{2(k+N-1)}).
$$

Here we have set the bosonic operators $\widetilde{S}_i(z)$ $(1 \leq i \leq N)$ by

$$\begin{aligned}
\widetilde{S}_i(z) &= \frac{1}{(q-q^{-1})z} \sum_{j=i+1}^{N} : \left(e^{-b_-^{i,j}(q^{N-1-j}z)-(b+c)^{i,j}(q^{N-j}z)} - e^{-b_+^{i,j}(q^{N-1-j}z)-(b+c)^{i,j}(q^{N-j-2}z)} \right) \\
&\times \ e^{(b+c)^{i+1,j}(q^{N-1-j}z)+\sum_{l=j+1}^{N}(b_-^{i+1,l}(q^{N-l}z)-b_-^{i,l}(q^{N-l-1}z))+b_-^{i+1,N+1}(z)-b_-^{i,N+1}(q^{-1}z)} : \\
&+ \ q : e^{b^{i,N+1}(z)+b_+^{i+1,N+1}(z)-b^{i+1,N+1}(qz)} : \qquad (1 \leq i \leq N-1), \\
\widetilde{S}_N(z) &= -q^{-1} : e^{b^{N,N+1}(z)} : .
\end{aligned}$$

The screening $\mathcal{Q}_i$ commutes with the quantum superalgebra.

$$[\mathcal{Q}_i, U_q(\widehat{sl}(N|1))] = 0 \quad (1 \leq i \leq N).$$

For $p_a^i \in \mathbf{C}$ $(1 \leq i \leq N)$, $p_b^{i,j} \in \mathbf{C}$ $(1 \leq i < j \leq N+1)$, $p_c^{i,j} \in \mathbf{C}$ $(1 \leq i < j \leq N)$, we set the vector $|p_a, p_b, p_c\rangle$ which satisfies the following conditions.

$$\begin{aligned}
&a_m^i|p_a,p_b,p_c\rangle = b_m^{i,j}|p_a,p_b,p_c\rangle = c_m^{i,j}|p_a,p_b,p_c\rangle = 0 \quad (m>0), \\
&a_0^i|p_a,p_b,p_c\rangle = p_a^i|p_a,p_b,p_c\rangle, \ b_0^{i,j}|p_a,p_b,p_c\rangle = p_b^{i,j}|p_a,p_b,p_c\rangle, \ c_0^{i,j}|p_a,p_b,p_c\rangle = p_c^{i,j}|p_a,p_b,p_c\rangle.
\end{aligned}$$

The boson Fock space $F(p_a,p_b,p_c)$ is generated by the bosons $a_m^i, b_m^{i,j}, c_m^{i,j}$ on the vector $|p_a,p_b,p_c\rangle$. We set the space $F(p_a)$ by

$$F(p_a) = \bigoplus_{\substack{p_b^{i,j}=-p_c^{i,j}\in\mathbf{Z} \ (1\leq i<j\leq N) \\ p_b^{i,N+1}\in\mathbf{Z} \ (1\leq i\leq N)}} F(p_a,p_b,p_c).$$

We would like to construct the vertex operator $\Phi^*(z)$ [10] which gives the intertwiner among $F(p_a)$ and the typical representation [13].

$$\Phi^*(z) : F(p_a) \longrightarrow F(p_a+l_a+x_a) \otimes V_z^{*S}, \quad \Phi^*(z) = \sum_j \Phi_j^*(z) \otimes v_j^*.$$

Here V_z^{*S} is the dual evaluation representation of the typical representation with the weight $l_a = (l_a^1, l_a^2, \cdots, l_a^N)$ [13]. The basis of V^{*S} is given by $\{v_j^*\}$. The coefficients are linear maps : $\Phi_j^*(z)$: $F(p_a+l_a+x_a) \longrightarrow F(p_a)$. For $l_a = (l_a^1, l_a^2, \cdots, l_a^N)$ and $x_a = (x_a^1, x_a^2, \cdots, x_a^N)$, we set the highest element $\Phi_1^*(z)$ of the vertex operator by

$$\Phi_1^*(z) =: \prod_{j=1}^{N} \mathcal{Q}_j^{x_j} \ \ e^{-\sum_{i,j=1}^{N}\left(\frac{l_a^i}{k+N-1}\frac{\mathrm{Min}(i,j)}{N-1}\frac{N-1-\mathrm{Max}(i,j)}{1}a^j\right)(q^k z|-\frac{k-N+1}{2})} :,$$

where $(x_1, x_2, \cdots, x_N)$ is related to x_a^i by $x_a^i = \sum_{j=1}^{N} A_{i,j} x_j$. Other elements $\Phi_j^*(z)$ $(j \neq 1)$ of the vertex operators are determined by the intertwining property. We conjecture that this bosonization of the vertex operator $\Phi^*(z)$ gives the intertwiner among our bosonization and the typical representation. Using the Gelfand-Zetlin basis [13], we have checked this conjecture in some cases for $N = 2, 3, 4$ [10]. We balance the "background charge" of the vertex operators by using the screening $\mathcal{Q}_i$. Sometimes we have to multiply nontrivial product of the screenings $\mathcal{Q}_i$ inside the vertex operator in order to have non-zero correlation functions (trace of vertex operators). At the end of this paper, we would like to give some comments on relating works [8, 12]. In [8] we study how to get the Wakimoto realization from our bosonization by ξ-η system. In [12] we study how to construct the elliptic deformed algebra $U_{q,p}(\widehat{sl}(M|N))$ from the quantum group $U_q(\widehat{sl}(M|N))$. Using deformation method developed in [12] we obtain a bosonization of the elliptic deformed algebra $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k.

Acknowledgement

This work is supported by the Grant-in-Aid for Scientific Research **C** (21540228) from Japan Society for Promotion of Science.

References

[1] Zhang F and Rice T 1988 *Phys.Rev.***B37** 3759

[2] Kac V 1977 *Adv.Math.***26**(1) 8

[3] Yamane H 1999 *Publ.Res.Inst.Math.Sci.* **35** 321

[4] Wakimoto M 1986 *Commun.Math.Phys.***104** 605

[5] Awata H, Odake S and Shiraishi J 1994 *Commun.Math.Phys.* **162** 61

[6] Awata H, Odake S and Shiraishi J 1997 *Lett. Math. Phys.***42** 271

[7] Kojima T 2011 *J.Math.Phys.***53** 013515.

[8] Kojima T 2011 The q-Wakimoto realization of the superalgebras $U_{q,p}(\hat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$ to appear in *Springer Proceedings : Statistics and Mathematics* (*Preprint* arXiv.1201.4222)

[9] Zhang Y and Gould M 2000 *J.Math.Phys.***41** 5577

[10] Kojima T 2012 *J.Math.Phys.***53** 083503

[11] Jimbo M, Konno H, Odake S and Shiraishi J 1999 *Commun.Math.Phys.***199** 605

[12] Kojima T 2011 *J.Phys.***A44** 485205

[13] Palev T and Tolstoy V 1991 *Commun.Math.Phys.***141** 549